

Nutrition and Lower Extremity Ulcers: Causality and/or Treatment

The International Journal of Lower
Extremity Wounds
2017, Vol. 16(4) 238–243
© The Author(s) 2017
Reprints and permissions:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/1534734617737639
journals.sagepub.com/home/ijl



Lauren Haughey, PT, MPT, WCC¹, and Adrian Barbul, MD, FACS^{1,2}

Abstract

The association between malnutrition, impaired wound healing, and the presence of chronic wounds has been recognized for a long time. It is widely believed that the lack of adequate nutrition increases the risk of developing wounds which have a great likelihood of progressing to chronicity due to lack of appropriate healing responses. This risk is particularly high in the aging population. For the individual patient, as well as patient populations, the diagnosis of malnutrition has been in dispute; further, there is lack of agreement of when and how to intervene nutritionally in those with wounds or healing deficits. This article examines the relationship of nutritional status with the presence and clinical evolution of leg ulcers in humans, focusing on diabetic and venous leg ulcers; we will further review the effect of nutritional therapy on the outcome of these ulcers.

Keywords

nutritional, healing deficits, malnutrition

Definition of Malnutrition

The World Health Organization defines malnutrition as “deficiencies, excesses or imbalances in a person’s intake of energy and/or nutrients”¹; Merriam Webster’s defines it “as faulty nutrition due to inadequate or imbalanced intake, impaired assimilation or utilization.”² In the United States, deficiency of adequate, nutritionally appropriate caloric intake has been the generally accepted definition, regardless of etiology (poor intake, defective absorption, lack of interest, obesity, etc). Energy or protein deficiency, Kwashiorkor or Marasmus, respectively, are seldom seen in pure forms in the Western world, with most malnourished patients or elderly displaying characteristics of both. A more recent study characterized malnutrition as the presence of low body mass index, unintentional weight loss, low serum albumin and food intake less than 60% of the daily estimated requirement.³ 40% to 60% of hospitalized older adults, up to 85% of nursing home residents and upward of 60% of home care patients are either in a state of malnutrition or at risk.⁴ Level of independence in the elderly both cognitively and physically are inversely associated with staggeringly high rates of malnutrition.⁵ Loss of appetite, financial distress and depression are further associated with the presence of malnutrition.⁵⁻⁷ Obesity as a phenomena of impaired nutritional status is a relatively new problem in the world, seemingly ever increasing primarily in the Western world. It is of note that the most recent statistics from the World Health Organization indicate approximately 462 million individuals

are underweight versus the nearly 1.9 billion adults classified as overweight or obese.¹

Assessment of Malnutrition

Comprehensive evaluation of nutritional status should include patient history, complete physical exam, anthropometric measurements, and laboratory analyses. Anthropometric measures used include height, weight, body mass index (BMI), and estimates of body fat and muscle mass. Recommended laboratory measurements include albumin, retinal-binding prealbumin, transferrin, complete blood count, serum folate, vitamin B₁₂, and cholesterol.⁸ Neither the physical nor biochemical assessments have not been found to correlate well with malnutrition and are time consuming, expensive, and often prone lack of reproducibility.

The nutritional assessment of the elderly is rendered more difficult because many of the clinical signs of malnutrition are also associated with normal aging. Various tools have been promulgated for the nutritional assessment of

¹Tennessee Valley VA Hospital, Nashville, TN, USA

²Vanderbilt University School of Medicine, Nashville, TN, Vanderbilt University School of Medicine, Nashville, TN, USA

Corresponding Author:

Adrian Barbul, Department of Surgery, Vanderbilt University School of Medicine, 1161 21st Avenue South D-5203 Medical Center North, Nashville, TN, USA.

Email: adrian.barbul@vanderbilt.edu

geriatric populations, aiming for simplicity, accuracy of prediction and reproducibility. These include (1) Nutritional Screening Initiative, levels I and II, which is well suited to community screening; it incorporates a simple nomogram to determine BMI and a questionnaire of clinical characteristics, eating habits, living environment, and mental and cognitive states plus laboratory data (albumin and serum cholesterol); (2) Mini Nutritional Assessment: created for the elderly, it can be applied in the ambulatory or hospital setting; it defines the condition of the malnourished patient, the risk of malnutrition, and the general nutritional status; it can be performed in approximately 10 minutes and is highly reproducible⁵; (3) Mini Nutritional Assessment–Short Form: created to reduce the time of administration to 10 minutes without losing diagnostic power, it correlates well the mini nutritional assessment, it has adequate sensitivity and specificity, and good internal consistence⁹; (4) Malnutrition Universal Screening Tool: initially, developed for non-institutionalized individuals, its use has been validated in various environments including hospital, clinics, and home residences¹⁰; drawbacks include the lack of any measure of functionality and a focus on acute disease; and (5) Subjective Global Assessment, designed to estimate nutritional status through clinical history and physical exploration, it has greater sensitivity and specificity; can be used to determine which patients require nutritional intervention and which would benefit from intensive nutritional support.¹¹

In patients with malnutrition, evaluation of the potential risk for wound development and/or healing becomes the next challenge. Tools that factor malnutrition/nutritional status and the development of wounds such as the Norton, Waterlow, and Braden scales are used routinely in the hospital/institutional settings.^{12,13} All the aforementioned tools address nutrition in addition to other key factors such as mobility/activity status, skin status, and moisture exposure. Clearly, development of wounds is multifactorial and nutritional deficits are a major contributor to risk for wound development. BMI alone is not a good predictor of protein deficiency. The National Pressure Ulcer Advisory Panel (NPUAP) evolved their own risk assessment that ranks mobility status along with baseline skin status at the base of the pyramid while perfusion, oxygenation, and nutritional status are on the second tier and age, health status, sensory perception and hematological measures are at the apex of the pyramid. Inadequate protein intake leading to edema in peripheral vasculature,¹⁴ and acute and chronic inflammation¹⁵ have all been implicated mechanistically in the evolution of wounds in the background of malnutrition.

Malnutrition and Lower Leg Ulcers

In examining the nexus between nutrition and lower leg chronic wounds, we will refer mainly to human studies. Additionally, where information exists in some types of wounds (acute, pressure, etc), we would assume that biological

principles apply generally to all wounds, particularly where specific data is not available on lower extremity wounds.

Lack of adequate nutrition impairs the wound collagen accumulation, impairs wound healing and promotes chronicity.¹⁶ Intracellular collagen synthesis has an absolute requirement for energy, amino acids, oxygen, trace minerals, and vitamins.¹⁷ The malnourished organism, which is in energy conserving mode and carries out minimal synthetic processes during acute injury phase, has to overcome this metabolic impediment for wound healing, which is above all a synthetic process

Chronic wounds are closely associated with malnutrition. This association is strongest with pressure ulcers, but not all studies agree.^{18,19} Pressure ulcers of the lower extremities are typically seen in the heels as well as the lateral malleoli; with lack of mobility in the aged population, wounds of the lower legs can be in numerous places. Pressure ulcers developed as the individual is unable to shift pressure off of bony prominence or devices (catheter tubing, etc). Although poor nutrition alone is not a strong risk factor for developing venous leg ulcers (VLUs), patients with VLU often have poor nutritional indices. For example, older patients (>70 years of age) with VLU have a higher incidence of malnutrition, as assessed by anthropometric indices (BMI, arm/calf circumference, history of weight loss) and dietary history/assessment.²⁰ A thorough study of a small population of obese patients with VLU noted similar findings.²¹ Total serum protein and albumin levels were below normal in 27% of patients with VLU.²² Dietary profiling of VLU patients reveals increased consumption of fat, sodium and sugar along with a decreased vitamin C intake, when compared with recommended daily allowance (RDA) recommendations.²³ Significant deficits in vitamins A and E, carotenes, and zinc have also been described in the setting of chronic VLU.²⁴

Patients with diabetic foot ulcers (DFUs) have similar or even higher degrees of malnutrition and nutritional deficits.²⁵ One-third of subjects older than 60 years with Wagner grade I to II ulcers of at least 4 weeks duration were found to have protein-calorie malnutrition.²⁶ Poor scores on Mini Nutritional Assessment are an independent predictor of amputation outcomes.²⁷ Furthermore, a BMI less than 25 kg/m² places diabetic patients at a higher risk for amputation. Nutritional status of diabetic patients correlates with the severity of the ulcer, risk of infection and with outcome; the severity of the DFU increases as the nutritional state deteriorates.²⁶

Nutritional Therapy and Lower Leg Ulcers

Given the close correlation between nutritional status and the occurrence of lower extremity ulcers, it is an easy for clinicians to advise that nutritional therapy should be part and parcel of any treatment plan for such patients. Although

advice for nutritional intervention is almost universal, there is paucity of data or agreement in multiple areas: how to administer nutrition (enteral, parenteral, topical), what to administer, role of special nutrients, length of therapy, end points of therapy, and so on. The matter gets even more confusing as there is some data regarding efficacy of nutritional intervention in pressure ulcers, but precious little strong evidence in lower limb ulcerations.

As mentioned before, successful wound healing requires metabolically the presence of calories, protein, trace minerals, vitamins, and oxygen. Goodson and Hunt²⁸ have pointed out that successful wound nutrition is host nutrition, meaning correction of host nutritional deficits is key to successful wound healing.²⁸ Thus, provision of adequate calories and protein is key to reversing nutritional deficits, which, as mentioned, are often present in patient with chronic ulcers. Depending on the level of stress and preexisting degree of malnutrition, intake of 30 to 35 kcal/kg body weight and 1.2 to 1.5 g protein/kg is recommended in order to correct and maintain nutritional health in most individuals with wounds.²⁹ Although this indication has been put forth for pressure ulcers, there is no reason not to believe that it would apply to patients with other chronic wounds and with nutritional deficits. The recommendation is to continue this level of caloric and protein intake until wounds are resolved.²⁹ Nutritional supplementation was found to be beneficial for venous stasis wounds but not for diabetic and sickle cell ulcers.³⁰

The nutraceutical role of single nutrients on collagen synthesis and/or wound healing is still being studied. We will review some of these below; most of the available data are not derived from study of lower extremity ulcers, but whatever biologic value to wound healing is demonstrated may also apply to all manners of wounds.

Amino Acids and Wound Healing

Amino acids are important for wound healing to occur. In the past three decades there has been increased recognition that several amino acids have specific effect of increasing collagen synthesis, in doses above and beyond those required for nutritional needs.

Arginine is considered an essential amino acid under periods of severe stress. In the growing animal much of the dietary arginine is used for the synthesis of connective tissue proteins and in the injured animal an increase in arginine intake would be expected to result in more reparative connective tissue synthesis.³¹

Using a micro-model where collagen accumulation occurs in a subcutaneously placed segments of polytetrafluoroethylene (PTFE) tubing, young healthy human volunteers (25-35 years) were found to have a significant increase in wound collagen deposition following oral supplementation with either 30 g of arginine aspartate (17 g of

free arginine) or 30 g of arginine HCl (24.8 g of free arginine) daily for 14 days.³² In a subsequent study of healthy elderly humans (67-82 years) daily supplements of 30 g of arginine aspartate for fourteen days resulted in significantly enhanced collagen and total protein deposition at the wound site when compared with placebo controls.³³ It should be noted that the effect of arginine is on collagen synthesis, as neither oral nor parenteral arginine affects epithelial healing.³²⁻³⁴

A small diverse group of patients with pressure ulcer treated with supplemental arginine, vitamin C, and zinc showed improved wound healing.³⁵ Similarly diets high in protein and arginine content led to better healing than standard hospital diet.³⁶ Markedly improved collagen synthesis was noted in healthy elderly humans given dietary supplementation of arginine, glutamine, and β -hydroxy β methylbutyrate (HMB).³⁷ A metabolite of leucine, HMB increases muscle mass and is most effective when coupled with exercise. In diabetic foot ulcer patients, supplemental arginine, glutamine, and HMB had a significant healing effect only in subjects with plasma albumin levels less than 40 g/L and in those with decreased limb perfusion, as measured by ankle-brachial indices.³⁸

Ornithine, a metabolite of arginine in the urea cycle, as well as ornithine keto-glutarate also enhance wound collagen deposition in animals, although no studies in humans have yet been carried out.^{39,40}

Wound Healing and Fatty Acids

Lipids, the primary structure of all cell membranes, play an integral though poorly understood role in the wound healing process. As a source of calories, other than glucose and protein, a diet replete in lipids may theoretically limit protein catabolism in the stressed or wounded state. Impaired wound healing has been documented in humans with essential fatty acid deficiencies.⁴¹

The role of fats in wound healing has not been widely studied. More recently interest in defining possible benefits of specific lipid types has emerged. The omega-3 fatty acids have anti-inflammatory properties by inhibiting eicosanoid production and other mediators such as platelet-activating factor (PAF), interleukin-1 (IL-1), and tumor necrosis factor- α (TNF- α).⁴²⁻⁴⁴ Animals consuming diets enriched with omega-3 fatty acids had weaker wounds 30 days after injury. The weaker wounds did not contain less collagen; rather the omega-3 supplementation was thought to impair the quality, cross-linking, or spatial orientation of collagen fibrils.⁴⁵ In a human suction blister model, the daily consumption of omega 3-fatty acids (given as supplemental fish oil that contained 26 times and 12 times the daily recommended dosage, respectively, for eicosapentaenoic acid [EPA] and docosahexaenoic acid [DHA]) raised blister fluid IL-1 levels and lengthened healing time.⁴⁶ It does appear that fish oil supplements, especially in large doses,

adversely affect the healing response; it should be borne in mind that omega fatty acids are stored in fatty tissue and can reach supranormal levels quickly.

Vitamins and Wound Healing

Vitamin C deficiency has historical significance to wound healing because of its relation to scurvy, which has as its central element a failure in collagen synthesis and cross-linking. Ascorbic acid is a specific co-substrate for the enzymes 4-prolyl hydroxylase and lysyl hydroxylase; biochemically, it is a reducing agent required for the conversion of proline and lysine to hydroxyproline and hydroxylysine, which are critical to collagen cross-linking and mechanical strength.⁴⁷

Although the recommended dietary allowance for vitamin C is 70-90 mg/d, the clinical spectrum of its administration varies widely, with doses as high as 5g/day being safely administered.⁴⁸ There is no evidence to suggest that massive doses of ascorbic acid are of any substantial benefit to wound healing in nondeficient states; conversely, there is no evidence that excess vitamin C is toxic.^{48,49}

Vitamin A deficiency impairs wound healing. Subsequently, vitamin A was shown to enhance wound healing even in nondeficient states. More important, vitamin A, administered either topically or systemically, reverses the anti-inflammatory effects of corticosteroids on wound healing.^{50,51} Vitamin A has also been proposed as therapy for wound healing impaired by diabetes, tumor formation, chemotherapeutic agents, or radiation. Vitamin A increases the inflammatory response in wounds through enhanced lysosomal membrane lability, increased macrophage influx and activation and stimulation of collagen synthesis. In the severely injured (such as burn victims), doses of vitamin A as high as 25 000 IU/d (5 times the recommended daily dose) have been advocated and used without any significant side effects. Larger doses of vitamin A do not improve wound healing further and prolonged excessive intake can be toxic; therefore, it is typically supplemented for 10 to 14 days to avoid any negative consequences.

Micronutrients

Of the numerous trace elements present in the body, copper, zinc, and iron have the closest relationship to wound healing. Copper is a required cofactor for cytochrome oxidase and the cytosolic antioxidant superoxide dismutase. Lysyl oxidase is a key copper enzyme utilized in the synthesis of connective tissue, where it catalyzes the crosslinking of collagen and strengthens the collagen framework.

Zinc, the most well-known element for wound healing, has been used empirically in dermatological conditions for centuries. Zinc is essential to wound healing in both animals and humans. It is a cofactor for RNA and DNA polymerase

and consequently involved in DNA synthesis, protein synthesis, and cellular proliferation. Zinc deficiency impairs the critical roles each of these processes play in wound healing. Zinc levels less than 100 µg/100 mL have been associated with decreased fibroblast proliferation and collagen synthesis.⁵² These defects are readily reversed with repletion of zinc to normal levels. Zinc levels can be depleted in settings of severe stress as well as in patients receiving long-term steroids. In these settings, it is recommended that patients receive both vitamin A and zinc supplements to improve wound healing. The current RDA for zinc is 15 mg. A randomized double-blind placebo controlled study in diabetic foot wounds (Wagner grades II, III, and IV) noted decreased measurements of length and width as well as significant changes in metabolic profiles with use of 220 mg zinc sulfate (50 mg elemental zinc) over a 12-week period.⁵³ It should be stressed that no studies have demonstrated improvement in wound healing after the administration of zinc to patients who are not zinc deficient.⁵⁴ Topical application of zinc in VLU can be of benefit when used in conjunction with appropriate wound bed preparation.⁵⁵

Iron is required for the hydroxylation of proline and lysine and as a result, severe iron deficiency can result in impaired collagen production. As a part of the oxygen transport system, iron can affect wound healing, but again, this only occurs in settings of severe iron-deficiency anemia. In the clinical setting, iron deficiency is quite common and can result from blood loss, infectious etiologies, malnutrition, or an underlying hematopoietic disorder. Unlike other deficiencies of trace elements, iron deficiency can be easily detected and treated. Hypovolemia rather than specific levels of hemoglobin have been shown to adversely affect wound healing most likely by decreasing oxygen and nutrient delivery to the healing wound site.

Other Nutrients and Wound Healing

Honey, ancient in use for wound care from ancient Egypt and Rome to warfare care through the centuries, has been "rediscovered" within the past 2 decades. In burns, ulcers, and mixed wounds, honey appears to have favorable wound healing properties, particularly in mild to moderate superficial burns.^{56,57} The high content in sugar, water, proteins, as well as the presence of some B complex vitamins, minerals antioxidants (flavonoids, ascorbic acid, catalase, selenium) and enzymes in honey (invertase, amylase, glucose oxidase) may explain the purported benefits as topical treatment, rather than as a nutrient. Honey exposed to gamma radiation is considered medical grade and can dilute wound exudate and activate neutrophils. Second, flavonoids have antibacterial activity. Third, honey has low free water activity preventing bacterial growth and drawing fluid away from the wound via osmosis. Last, the pH of honey inhibits growth of microorganisms. Application of honey or honey

dressings to VLU has not improved outcomes at 12 weeks.⁵⁸ Comparison of honey application on DFUs with povidone iodine did not find a significant effect in healing rates with the use of honey.⁵⁹ All in all, the support for wide topical usage of honey on wounds remains unproven.⁶⁰

The nexus of nutrition to wounds and wound healing is very strong. There is a paucity of high-quality data regarding the use of nutrition in reversing wound risk or improving wound healing. Recommendations to add 500 mg vitamin C, 17 mg zinc, and 17 g arginine daily have been made and soft data provided for such use; additionally, it appears totally safe. Prevention of wounds is multifaceted, reversing malnutrition is a monumental task and treatment of wounds as they relate to one's state of nutrition continues to be a major clinical challenge.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

References

- World Health Organisation. Malnutrition: fact sheet. <http://www.who.int/mediacentre/factsheets/malnutrition/en/>. Accessed October 3, 2017.
- Merriam-Webster. Malnutrition. <https://www.merriam-webster.com/dictionary/malnutrition#medicaldictionary>. Accessed October 3, 2017.
- Cereda E, Pedrolli C, Klersy C, et al. Nutritional status in older persons according to healthcare setting: a systematic review and meta-analysis of prevalence data using MNA®. *Clin Nutr*. 2016;35:1282-1290.
- Langemo D, Anderson J, Hanson D, Hunter S, Thompson P, Posthauer M. Nutritional considerations in wound care. *Adv Skin Wound Care*. 2006;19:297-298,300,303.
- Guigoz Y, Lauque S, Vellas BJ. Identifying the elderly at risk for malnutrition the Mini Nutritional Assessment. *Clin Geriatr Med*. 2002;18:737-757.
- Woo J, Lynn H, Lau WY, et al. Nutrient intake and psychological health in an elderly Chinese population. *Int J Geriatric Psychiatry*. 2006;21:1036-1043.
- Brownie S. Why the elderly are at risk for nutritional deficiency? *Intl J Nurs Pract*. 2016;12:110-118.
- Juárez-Cedillo T. Changes in nutritional needs with aging. In: Watson RR, ed. *Nutritional and Functional Foods for Healthy Aging*. San Diego, CA: Academic Press; 2017:27-41.
- Rubenstein LZ, Harker JO, Salvà A, Guigoz Y, Vellas B. Screening for undernutrition in geriatric practice: developing the short-form mini-nutritional assessment (MNA-SF). *J Gerontol A Biol Sci Med Sci*. 2001;56:M366-M372.
- Stratton RJ, Hackston A, Longmore D, et al. Malnutrition in hospital outpatients and inpatients: prevalence, concurrent validity and ease of use of the "malnutrition universal screening tool" (MUST) for adults. *Br J Nutr*. 2004;92:799-808.
- Detsky AS, McLaughlin JR, Baker JP, et al. What is the subjective global assessment of nutritional status? *JPEN J Parenter Enteral Nutr*. 1987;11:8-13.
- González-Ruiz JM, Sebastián-Viana T, Losa-Iglesias ME, et al. Braden scale and Norton scale modified by INSALUD in an acute care hospital: validity and cutoff point. *Adv Skin Wound Care*. 2014;27:506-511.
- Pancorbo-Hidalgo PL, Garcia-Fernandez FP, Lopez-Medina IM, Alvarez-Nieto C. Risk assessment scales for pressure ulcer prevention: a systematic review. *J Adv Nurs*. 2006;54:94-110.
- Randall HT. Water, electrolytes and acid-base balance. In Goodhart RS, Shils M, eds. *Modern Nutrition in Health and Disease*. 6th ed. Philadelphia, PA: Lea & Febiger; 1980:324-361.
- Jensen GL, Mirtallo J, Compher C, et al. Adult starvation and disease-related malnutrition: a proposal for etiology-based diagnosis in the clinical practice setting from the International Consensus Guideline Committee. *Clin Nutr*. 2010;29:151-153.
- Windsor JA, Knight GS, Hill GL. Wound healing response in surgical patients: recent food intake is more important than nutritional status. *Br J Surg*. 1988;75:135-137.
- Barbul A, Efron DT, Kavalukas SL. Wound healing. In: Brunicaardi FC, Andersen DK, Billiar TR, et al, eds. *Schwartz's Principles of Surgery*. 10th ed. New York, NY: McGraw Hill; 2014:241-272.
- Posthauer ME, Banks M, Dorner B, Schols JM. The role of nutrition for pressure ulcer management: National Pressure Ulcer Advisory Panel, European Pressure Ulcer Advisory Panel, and Pan Pacific Pressure Injury Alliance white paper. *Adv Skin Wound Care*. 2015;28:175-188.
- Langer G, Fink A. Nutritional interventions for preventing and treating pressure ulcers. *Cochrane Database Syst Rev*. 2014;(6):CD003216.
- Szewczyk MT, Jawien A, Kedziora-Kornatowska K, et al. The nutritional status of older adults with and without venous ulcers: a comparative, descriptive study. *Ostomy Wound Manage*. 2008;54:34-36, 38-40, 42.
- Tobón J, Whitney JD, Jarrett M. Nutritional status and wound severity of overweight and obese patients with venous leg ulcers: a pilot study. *J Vasc Nurs*. 2008;26:43-52.
- Legendre C, Debure C, Meaume S, Lok C, Golmard JL, Senet P. Impact of protein deficiency on venous ulcer healing. *J Vasc Surg*. 2008;48:688-693.
- McDaniel JC, Kemmner KG, Rusnak S. Nutritional profile of older adults with chronic venous leg ulcers: a pilot study. *Geriatr Nurs*. 2015;36:381-386.
- Rojas AL, Phillips TJ. Patients with chronic leg ulcers show diminished levels of vitamins A and E, carotenes and zinc. *Dermatol Surg*. 1999;25:601-604.
- Zhang SS, Tang ZY, Fang P, Qian HJ, Xu L, Ning G. Nutritional status deteriorates as the severity of diabetic foot

- ulcers increases and independently associates with prognosis. *Exp Ther Med*. 2013;5:215-222.
26. Gau BR, Chen HY, Hung SY, et al. The impact of nutritional status on treatment outcomes of patients with limb-threatening diabetic foot ulcers. *J Diabetes Complications*. 2016;30:138-142.
 27. Yetka Z, Pourali R, Nezhadrahim R, Ravanyar L, Ghasemi-Rad M. Clinical and behavioral factors associated with management outcomes in hospitalized patients with diabetic foot ulcers. *Diabetes Metab Syndr Obes*. 2011;4:371-375.
 28. Goodson WH 3rd, Hunt TK. Wound Healing and Nutrition. In: Kinney JM, Jeejeebhoy KN, Hill GL, Owen OE, eds. *Nutrition and Metabolism in Patient Care*. Philadelphia, PA: W B Saunders; 1988:635-642.
 29. National Pressure Ulcer Advisory Panel; European Pressure Ulcer Advisory Panel; Pan Pacific Pressure Injury Alliance; National Guideline Clearinghouse. *Prevention and Treatment of Pressure Ulcers; Clinical Practice Guideline*. Washington, DC: National Pressure Ulcer Advisory Panel; 2009.
 30. Ye J, Mani R. A systematic review and meta-analysis of nutritional supplementation in chronic lower extremity wounds. *Intl J Low Extrem Wounds*. 2016;15:296-302.
 31. Barbul A. Arginine: biochemistry, physiology, and therapeutic implications. *JPEN J Parenter Enteral Nutr*. 1986;10:227-238.
 32. Barbul A, Lazarou SA, Efron DT, Wasserkrug HL, Efron G. Arginine enhances wound healing in humans. *Surgery*. 1990;108:331-337.
 33. Kirk SJ, Hurson M, Regan MC, Holt DR, Wasserkrug HL, Barbul A. Arginine stimulates wound healing and immune function in aged humans. *Surgery*. 1993;114:155-160.
 34. Debats IB, Wolfs TG, Gotoh T, Cleutjens JP, Peutz-Kootstra CJ, van der Hulst RR. Role of arginine in superficial wound healing in man. *Nitric Oxide*. 2009;21:175-183.
 35. Desneves KJ, Todorovic BE, Cassar A, Crowe TC. Treatment with supplemental arginine, vitamin C, and zinc in patients with pressure ulcers: a randomized controlled trial. *Clin Nutr*. 2005;24:979-987.
 36. Benati G, Delvecchio S, Cilla D, Pedone V. Impact of pressure ulcer healing of an arginine enriched solution in patients with severe cognitive impairment. *Arch Gerontol Geriatr Suppl*. 2001;7:43-47.
 37. Williams JZ, Abumrad NN, Barbul A. Effect of a specialized amino acid mixture on human collagen deposition. *Ann Surg*. 2002;236:369-375.
 38. Armstrong DG, Hanft JR, Driver VR, et al; Diabetic Food Nutrition Study Group. Effect of oral nutritional supplementation on wound healing in diabetic foot ulcers: a prospective randomized controlled trial. *Diabet Med*. 2014;31:1069-1077.
 39. Shi HP, Fishel RS, Efron DT, Williams JZ, Fishel MH, Barbul A. Effect of supplemental ornithine on wound healing. *J Surg Res*. 2002;106:299-302.
 40. Uzgar A, Cynober L, Barbul A. Ornithine alpha ketoglutarate enhances wound healing. *J Am Coll Surg*. 2009;209:S72.
 41. O'Neill JA Jr, Caldwell MD, Meng HC. Essential fatty acid deficiency in surgical patients. *Ann Surg*. 1977;185:535-542.
 42. Kremer JM, Jubiz W, Michalek A, et al. Fish oil fatty acid supplementation in active rheumatoid arthritis. A double-blind, controlled, crossover study. *Ann Int Med*. 1987;106:497-503.
 43. Sperling RI, Robin JL, Kylander KA, Lee TH, Lewis RA, Austen KF. The effects of N-3 polyunsaturated fatty acids on the generation of platelet-activating factor by human monocytes. *J Immunol*. 1987;139:4186.
 44. Endres S, Ghorbani R, Kelley VE, et al. The effect of N-3 polyunsaturated fatty acids on the synthesis of interleukin-1 and tumor necrosis factor by mononuclear cells. *N Engl J Med*. 1989;320:265-271.
 45. Albina JE, Gladden P, Walsh WR. Detrimental effects of omega-3 fatty acid-enriched diet on wound healing. *J Parent Ent Nutr*. 1993;17:519-521.
 46. McDaniel JC, Belury M, Ahijevych K, Blakely W. Omega 3-fatty acids effect on wound healing. *Wound Repair Regen*. 2008;16:337-345.
 47. Kivirikko KI, Helaakoski T, Tesanen K, et al. Molecular biology of prolyl 4-hydroxylase. *Ann NY Acad Sci*. 1990;580:132-142.
 48. Schmidt KH, Hagmaier V, Hornig DH, Vuilleumier JP, Rutishauser G. Urinary oxalate excretion after large intakes of ascorbic acid in man. *Am J Clin Nutr*. 1981;34:305-311.
 49. Rivers JM. Safety of high-level vitamin C injection. *Ann NY Acad Sci*. 1987;498:445-454.
 50. Brandaleone H, Papper E. The effect of the local and oral administration of cod liver oil on the rate of wound healing in vitamin A-deficient and normal rats. *Ann Surg*. 1941;114:791-798.
 51. Ehrlich HP, Hunt TK. Effects of cortisone and vitamin A on wound healing. *Ann Surg*. 1968;167:324-328.
 52. Lansdown AB, Mirastschijski U, Stubbs N, Scanlon E, Agren MS. Zinc in wound healing: theoretical, experimental, and clinical aspects. *Wound Repair Regen*. 2007;15:2-16.
 53. Momen-Heravi M, Barahimi E, Razzaghi R, Bashmani F, Gilasi HR, Asemi Z. The effects of zinc supplementation on wound healing and metabolic status in patients with diabetic foot ulcer: a randomized, double-blind, placebo-controlled trial. *Wound Repair Regen*. 2017;25:512-520.
 54. Wilkinson EA. Oral zinc for arterial and venous leg ulcers. *Cochrane Database Syst Rev*. 2012;(8):CD001273.
 55. O'Connor S, Murphy S. Chronic venous ulcers: is topical zinc the answer? A review of the literature. *Adv Skin Wound Care*. 2014;27:35-44.
 56. Vandamme L, Heyneman, Hoeksema H, Verbelen J, Monstrey SM. Honey in modern wound care: a systematic review. *Burns*. 2013;39:1514-1525.
 57. Jull AB, Rodgers A, Walker N. Honey as topical treatment for wounds. *Cochrane Database Syst Rev*. 2008;(4):CD005083.
 58. Jull A, Walker N, Parag V, Molan P, Rodgers A; Honey as Adjuvant Leg Ulcer Therapy Trial Collaborators. Randomized clinical trial of honey-impregnated dressings for venous leg ulcers. *Br J Surg*. 2008;95:175-182.
 59. Shukrimi A, Sulaiman AR, Halim AY, Azril A. A comparative study between honey and povidine iodine as a dressing solution for Wagner II diabetic foot ulcers. *Med J Malaysia*. 2008;63:44-46.
 60. Jull AB, Walker N, Deshpande S. Honey as a topical treatment for wounds. *Cochrane database Syst Rev*. 2013;(2):CD005083. doi:10.1002/14651858.CD005083.pub3.